


The background of the slide features a series of thin, parallel diagonal lines that create a sense of depth and movement. The color gradient transitions from a warm orange-brown on the left to a cool blue on the right. At the very top and bottom edges, there are horizontal bars with a slight glow. A white arrow points to the right on the top bar, and another white arrow points to the left on the bottom bar.

Measuring thermal Hall effect in a chiral spin liquid

Leon Balents, KITP

Collaborators


- Mengxing Ye (Minnesota)



- Gábor Halász (KITP -> ORNL)



- Lucile Savary (ENS Lyon)



Quantum Spin Liquid



Phil Anderson, 1973

a “quantum liquid” of spins

$$\text{blue oval} = \frac{1}{\sqrt{2}} (|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle)$$


$$\Psi = \text{diagram 1} + \text{diagram 2} + \dots$$

The diagram shows two triangular lattices of blue ovals. In the first lattice, the ovals are arranged in a regular pattern. In the second lattice, the ovals are arranged in a different pattern, representing a different state. The two lattices are added together, followed by an ellipsis indicating further terms in the sum.

Resonating **V**alence **B**ond state

Fractional quantum number

spinon



excitation with $\Delta S = 1/2$
not possible for any finite
cluster of spins

always created in pairs by any
local operator

Smoking Gun?

Difficult to find incontrovertible and sharp indicator of a QSL!

A few possibilities

- T-linear thermal conductivity in spinon Fermi surface QSL **observed!**
- Transverse, linearly dispersing emergent photon mode in 3d U(1) QSL **not yet observed**




Kitaev model

Kitaev's honeycomb model

$$H = \sum_{i,\mu} K_{\mu} \sigma_i^{\mu} \sigma_{i+\mu}^{\mu}$$

1. The model



Spin $\frac{1}{2}$ on each site.

exact parton construction $\sigma_i^{\mu} = i c_i c_i^{\mu}$ $c_i c_i^x c_i^y c_i^z = 1$


effective quadratic Hamiltonian

$$H = \sum_{i,\mu} i K_{\mu} c_i c_{i+\mu}$$




Gapless QSL


$$H = \sum_{i,\mu} iK_{\mu} c_i c_{i+\mu}$$




~1/4 of graphene




Phase diagram



Non-local excitations



Majorana ε




Flux e, m

In Kitaev's model:

- Majorana's dispersion $\sim K$ and Dirac-like
- Fluxes are localized with small gap

Non-Abelian Phase

- In an applied magnetic field, the Majoranas acquire a gap




$$H_{\text{eff}} = \frac{i}{4} \sum_{j,k} A_{jk} c_j c_k,$$

$$A = 2J (\text{solid arrow}) + 2\kappa (\text{dashed arrow}),$$

$$\kappa \sim \frac{h_x h_y h_z}{J^2}.$$

field induces a fermion mass, very similar to the Haldane model (except Majorana)



$$H_e = -\frac{iv}{4} \int dx \eta \partial_x \eta$$

Quantum Hall Effect?

- No charge. Have to study heat transport!

T



$$I = \int_0^\infty \frac{dq}{2\pi} v^2 q f(vq) = \frac{c\pi k_B^2}{12\hbar} T^2$$


central charge $c=1/2$

c.f. $c=1$ for both IQHE and FQHE abelian states

implies the existence of bulk non-abelian excitations (the fluxes, bound to MZMs)

Quantum Hall Effect?


- No charge. Have to study heat transport!



$$I = \frac{c\pi k_B^2}{12\hbar} (T_1^2 - T_2^2)$$
$$\approx \frac{c\pi k_B^2 T}{6\hbar} (T_1 - T_2)$$

Quantum Hall Effect?

- No charge. Have to study heat transport!



$$I_x = \kappa_H \Delta T_y$$



$$\kappa_H = \frac{\pi c k_B^2 T}{6\hbar}$$

a universal prediction for chiral
"Ising anyon" phase: *agnostic to
microscopic spin interactions*


Kitaev Materials

Jackeli, Khaliullin
2009

Showed that Kitaev interaction can be large in edge-sharing octahedra with large spin-orbit-coupling



Na_2IrO_3 ,
 (α, β, γ) -
 Li_2IrO_3



$\alpha\text{-RuCl}_3$



Y.-J. Kim...

P.Gegenwart
H. Takagi
...

Honeycomb and hyper-honeycomb structures

$\alpha\text{-RuCl}_3$

Clearly not described by ideal Kitaev model,
but still interesting




BUT: excitations to $\sim 15\text{meV}$ = 170K

Suppressing order

With a magnetic field:

A. Banerjee et al, 2017




spin liquid or boring
paramagnet?


Suppressing order

With a magnetic field:

A. Banerjee et al, 2017





theoretical possibilities






Thermal Hall Effect

Y. Kasahara *et al*, Nature 2018






Thermal Hall Effect

Y. Kasahara *et al*, Nature 2018






Thermal Hall Effect

Y. Kasahara *et al*, Nature 2018




Thermal Hall Effect

Y. Kasahara *et al*, Nature 2018



Why worry?



c.f. IQHE

Quantization is *only* visible when diagonal conductivity is small, i.e. Hall angle $\sim 90^\circ$

semicircle "law"

$$\sigma_{xx}^2 + \left(\sigma_{xy} - \frac{\sigma_1 + \sigma_2}{2} \right)^2 = \left(\frac{\sigma_1 - \sigma_2}{2} \right)^2$$

But for RuCl_3 , thermal Hall angle $\sim 10^{-3} !!$

Theoretical Problem

From Y. Kasahara *et al*


In the plateau regime of κ_{xy} , no anomaly is observed in κ_{xx} , probably because **phonon contributions largely dominate** over fermionic excitations arising from spins in κ_{xx} in the whole temperature range^{28,29}.

Q: Can quantized thermal Hall effect persist when phonons dominate bulk conductivity?

A: **Yes**, and phonons actually *help* to make the effect observable.


Theoretical Problem

Recall
derivation of
QTHE




Theoretical Problem

Recall
derivation of
QTHE




- $\kappa_{xx} \gg \kappa_{xy}$: Temperature of phonons is *not* constant at the edge
?? Could edge be out of equilibrium with the phonons and have constant T ??
- but: in that case, which T is measured?

Formulation




- Assume spins go into nonabelian QSL phase: bulk spin gap, chiral edge state
- Include bulk phonons with *no* Hall conductivity
- Majoranas and phonons can exchange energy at the edge of the sample

Formulation



- Variables:
 - $T_{ph}(x,y)$ = phonon temperature in bulk
 - $T_f(x,L_y)$ etc = fermion temperature at edge

Formulation



leads coupled
to the lattice

- Currents:


- Bulk

$$\mathbf{j}_{ph} = -\kappa \nabla T_{ph}$$

- Edge

$$I_f = \frac{\pi}{24} T_f^2$$

Formulation




- Continuity:

- Bulk $\nabla \cdot \mathbf{j}_{ph} = 0$


- Edge $\partial_x I_f = \pm j_{ex}, \quad \pm j_{ph}^y(x, \pm y_0) = j_{ex}(x, \pm y_0).$

Formulation




- Coupling:
- linear response $j_{ex} = \lambda(T)(T_{ph} - T_f)$

Formulation



- Full equations (for *infinite* Hall bar)
- bulk $\nabla^2 T_{ph} = 0$
- edge $\kappa_{xy}^q \partial_x T_f(x, \pm y_0) = -\kappa \partial_y T_{ph}(x, \pm y_0)$
 $= \pm \lambda(T) (T_{ph} - T_f)|_{x, \pm y_0}$

Solution



- Solution


- Bulk $T_{ph}(x, y) = T_0 + \frac{\partial T_{ph}}{\partial x}x + \frac{\partial T_{ph}}{\partial y}y$

- edge $T_f(x, \pm y_0) = T_{1,\pm} + \frac{\partial T_{ph}}{\partial x}x$

$$\frac{\partial T_{ph}}{\partial y} = -\frac{\kappa_{xy}^q}{\kappa} \frac{\partial T_{ph}}{\partial x}$$

$$T_{ph} - T_f = \mp \frac{\kappa}{\lambda} \frac{\Delta T_y}{L_y}$$

Solution



- Solution

- Bulk $T_{ph}(x, y) = T_0 + \frac{\partial T_{ph}}{\partial x}x + \frac{\partial T_{ph}}{\partial y}y$


- edge $T_f(x, \pm y_0) = T_{1,\pm} + \frac{\partial T_{ph}}{\partial x}x$

$$\frac{\partial T_{ph}}{\partial y} = -\frac{\kappa_{xy}^q}{\kappa} \frac{\partial T_{ph}}{\partial x}$$

$$T_{ph} - T_f = \mp \frac{\kappa}{\lambda} \frac{\Delta T_y}{L_y}$$

phonon Hall angle corresponds to quantization! **Implies**
quantization in 3 probe experiment measuring T_{ph}

Solution



- Solution

- Bulk $T_{ph}(x, y) = T_0 + \frac{\partial T_{ph}}{\partial x}x + \frac{\partial T_{ph}}{\partial y}y$


- edge $T_f(x, \pm y_0) = T_{1,\pm} + \frac{\partial T_{ph}}{\partial x}x$

$$\frac{\partial T_{ph}}{\partial y} = -\frac{\kappa_{xy}^q}{\kappa} \frac{\partial T_{ph}}{\partial x}$$

$$T_{ph} - T_f = \mp \frac{\kappa}{\lambda} \frac{\Delta T_y}{L_y}$$


Majorana temperature *not* equal to phonon one. It is actually *better* to measure phonon temperature!

Finite Hall bar



- Solve coupled Laplace and boundary equations, with constant temperature leads.
- Reveals dependence on “thermalization length”

$$\ell_x = \frac{\kappa_{xy}^q}{\lambda}$$



$$L_y/L_x = 0.01$$

$$\ell_x/L_x = \underline{0.01}, \underline{0.1}, \underline{1}$$


Bulk-edge coupling

local phonon strain couples
to Majorana kinetic energy

$$H_{int} = \frac{-igv_f}{4} \int dx \zeta(x) K_{ij} \partial_i u_j(x, y_0) \eta(x) \partial_x \eta(x),$$

We use kinetic equation to
calculate

$$\left(\frac{\partial \mathcal{E}}{\partial t}\right)_{ph \rightarrow f} = -\left(\frac{\partial \mathcal{E}_{ph}}{\partial t}\right) = -\sum_{q_x, q_y} \omega_{\vec{q}} \left(\frac{\partial g(\vec{q})}{\partial t}\right)_{coll},$$



Rate

$$\begin{aligned} \left(\frac{\partial g(\vec{q})}{\partial t}\right)_{coll} = \frac{2\pi}{\hbar} \times 2 \sum_{k, k'} \{ & \langle |\mathcal{M}^+(\vec{q}, k, k')|^2 \rangle_{dis} (1 + g_\omega) f_\epsilon f_{\epsilon'} \delta(-\omega_{\vec{q}} + \epsilon_k + \epsilon_{k'}) \\ & - \langle |\mathcal{M}^-(\vec{q}, k, k')|^2 \rangle_{dis} g_\omega f_\epsilon f_{\epsilon'} \delta(\omega_{\vec{q}} + \epsilon_k + \epsilon_{k'}) \} \end{aligned}$$


Bulk-edge coupling

local phonon strain couples
to Majorana kinetic energy

$$H_{int} = \frac{-igv_f}{4} \int dx \zeta(x) K_{ij} \partial_i u_j(x, y_0) \eta(x) \partial_x \eta(x),$$

We use kinetic equation to

calculate $j_{ex} \propto \left(\frac{\partial \mathcal{E}}{\partial t} \right)_{ph \rightarrow f} = - \left(\frac{\partial \mathcal{E}_{ph}}{\partial t} \right) = - \sum_{q_x, q_y} \omega_{\vec{q}} \left(\frac{\partial g(\vec{q})}{\partial t} \right)_{coll},$




Result

$$\lambda = \frac{g^2 \zeta^2}{32(2\pi)^3 v_{ph}^4 v_f^2 \rho_0} f T^6$$

**strong T-dependence:
thermalization becomes
poor at low T.**

Prediction




Expect *failure* of quantization at low enough temperature!

Disorder

- Experimental observation:
 - Quantization is observed only in the “best” samples, which have the largest *diagonal* thermal conductivity
- Presumably, sample variations involve lattice disorder that affects the phonon transport
- What is the effect on thermal Hall effect?

Disorder

- Network model



— resistor $I_{i \rightarrow j} = \kappa_{ij}(T_i - T_j)$


→ chiral $I_{i \rightarrow j} = \pm \kappa_H(T_i + T_j)/2$

- - - resistor $I_{i \rightarrow j} = \lambda(T_i - T_j)$

- Disorder included as random resistors

Disorder

Input: smooth random conductivity




Output: Hall temperature gradient
(anti-symmetrized in field)




Disorder


Input: smooth random conductivity



Output: Hall temperature gradient
(anti-symmetrized in field)



Summary



- Quantized value of thermal Hall conductivity can be measured even when large phonon conductivity is present
- Quantization is only power-law good, and deviations are controlled by edge-bulk equilibration

ARXIV:1805.10532

see also: ARXIV:1805.11507



U.S. DEPARTMENT OF
ENERGY

Office of
Science

GORDON AND BETTY
MOORE
FOUNDATION