

© Louis Taillefer

© LB

© LB

Hunting spin liquids: checking out the terrain and identifying the tracks

Leon Balents, UCSB

KITP conference "Gapless24", May 2024

Collaborators



Zhenhao Song
UCSB



Zhu-Xi Luo
Harvard
→ Georgia Tech




Urban Seifert
U. Köln



Hongchen Jiang
SLAC




Oleg Starykh
U. Utah



Zi-Yang Meng
HKU


Quantum spin liquids

- Topological QSLs




TQFT

- U(1) QSL




4d Maxwell

- Dirac QSLs





QED3

- Spinon Fermi surface




QED3 with
 $\mu \neq 0$





Spin liquid terrain

Quantum simulator




219 qubits

Very high degree of control

athermal

2d materials



10^6 qubits

Some control

meV scales

Crystals




10^{22} qubits

Caveat emptor

eV scales

TMD





From wikipedia



M=transition metal,
W, Mo, Nb etc.

X=chalcogenide
S, Se, Te


Typical band structure of TMD



- Quadratic dispersion: non-relativistic Schrödinger equation
- Spin-valley locking

F. Wu et al, 2018

TMD




From wikipedia



M=transition metal,
W, Mo, Nb etc.

X=chalcogenide
S, Se, Te

Typical band structure of bilayer



	graphene	TMD
Typical energy scales	$W \approx 10\text{-}100\text{ meV}$ $U \approx 20\text{-}40\text{ meV}$	$W \approx 1\text{-}100\text{ meV}$ $U \approx 100\text{-}200\text{ meV}$

K.-F. Mak, J. Shan, 2022

Hubbard simulator

PHYSICAL REVIEW LETTERS **121**, 026402 (2018)

Hubbard Model Physics in Transition Metal Dichalcogenide Moiré Bands

Fengcheng Wu,¹ Timothy Lovorn,² Emanuel Tutuc,³ and A. H. MacDonald²

PHYSICAL REVIEW B **102**, 201104(R) (2020)

Quantum phase diagram of a Moiré-Hubbard model

Haining Pan¹, Fengcheng Wu², and Sankar Das Sarma

Article

Simulation of Hubbard model physics in WSe₂/WS₂ moiré superlattices

<https://doi.org/10.1038/s41586-020-2085-3> Yanhao Tang¹, Lizhong Li¹, Tingxin Li¹, Yang Xu¹, Song Liu², Katayun Barmak³, Kenji Watanabe⁴, Takashi Taniguchi⁴, Allan H. MacDonald⁵, Jie Shan^{1,6,7,8} & Kin Fai Mak^{1,6,7,8}
Received: 18 August 2019

PHYSICAL REVIEW B **104**, 075150 (2021)

Editors' Suggestion

Hartree-Fock study of the moiré Hubbard model for twisted bilayer transition metal dichalcogenides

Jiawei Zang¹, Jie Wang², Jennifer Cano^{2,3} and Andrew J. Millis^{1,2}

moiré (extended) Hubbard model

$$H = \sum_s \sum_{i,j} t_s(\mathbf{R}_i - \mathbf{R}_j) c_{i,s}^\dagger c_{j,s} + \frac{1}{2} \sum_{s,s'} \sum_{i,j} U(\mathbf{R}_i - \mathbf{R}_j) c_{i,s}^\dagger c_{j,s'}^\dagger c_{j,s'} c_{i,s},$$


complex hopping Longer-range interactions

Caution: does not apply if active band is topologically non-trivial

The Hubbard model is hard

D. Arovas et al,
ARCMP, 2022

~60 years!



c.f.


Simons Collaboration on the
Many Electron Problem

Figure 5

Speculative ground state phase diagrams of the Hubbard model as a function of U and chemical potential, μ


Moiré Hubbard simulator

WS₂/WSe₂ bilayers



Y. Xu et al, 2020


Fractional filling insulators: charge order



X. Huang et al, 2021


Emergent lattices

d $\nu = \frac{1}{3} \left(\frac{2}{3} \right)$



Triangular/honeycomb

e $\nu = \frac{1}{4} \left(\frac{3}{4} \right)$





Triangular/kagomé

The Wigner crystallization is essentially classical

What are the spins doing?

Emergent lattices


d $\nu = \frac{1}{3} \left(\frac{2}{3} \right)$



H. Li *et al*, 2022

Electron and hole excitations
in twisted WS₂ imaged via
graphene capping layer

e $\nu = \frac{1}{4} \left(\frac{3}{4} \right)$




Triangular/kagomé


The Wigner crystallization is essentially classical

What are the spins doing?

Hartree-Fock studies



J. Zang et al, 2021



+ many more

H. Pan + S. Das Sarma, 2022

These studies are incapable of finding spin liquids

(Much fewer studies with other techniques, not global)

Slave rotor

$$c_{i\alpha} = f_{i\alpha} b_i$$

$$f_i^\dagger f_i = n_i \quad \text{gauge constraint}$$

$$b_i = e^{i\varphi_i}$$

$$[n_i, \varphi_j] = i\delta_{ij}$$

S. Florens + A. Georges, 2004

Slave rotor

$$c_{i\alpha} = f_{i\alpha} b_i$$

$$f_i^\dagger f_i = n_i \quad \text{gauge constraint}$$

$$b_i = e^{i\varphi_i}$$

$$[n_i, \varphi_j] = i\delta_{ij}$$

S. Florens + A. Georges, 2004

Due to the constraint, density-density interactions can be cast entirely into the boson sector

$$H = \sum_{ij;\alpha} t_{ij,\alpha} f_{i\alpha}^\dagger f_{j\alpha} b_i^\dagger b_j + \frac{1}{2} \sum_{i,j} U_{ij} n_i n_j$$



$$H_{mf} = \sum_{ij;\alpha} t_{ij,\alpha}^{\text{eff}} f_i^\dagger f_j + K_{ij} b_i^\dagger b_j + \frac{1}{2} \sum_{i,j} U_{ij} n_i n_j$$

Slave rotor

$$c_{i\alpha} = f_{i\alpha} b_i \qquad f_i^\dagger f_i = n_i \quad \text{gauge constraint}$$

$$b_i = e^{i\varphi_i}$$

$$[n_i, \varphi_j] = i\delta_{ij}$$

Due to the constraint, density-density interactions can be cast entirely into the boson sector

$$\text{MF conditions} \quad t_{ij,\alpha}^{\text{eff}} = t_{ij,\alpha} \langle b_i^\dagger b_j \rangle \quad K_{ij} = \sum_{\alpha} t_{ij,\alpha} \langle f_{i,\alpha}^\dagger f_{j,\alpha} \rangle$$

$$H_{mf} = \sum_{ij;\alpha} t_{ij,\alpha}^{\text{eff}} f_i^\dagger f_j + K_{ij} b_i^\dagger b_j + \frac{1}{2} \sum_{i,j} U_{ij} n_i n_j$$

Slave rotor

$$c_{i\alpha} = f_{i\alpha} b_i \quad f_i^\dagger f_i = n_i \quad \text{gauge constraint}$$

$$b_i = e^{i\varphi_i}$$

$$[n_i, \varphi_j] = i\delta_{ij}$$

Due to the constraint, density-density interactions can be cast entirely into the boson sector

$$\text{MF conditions} \quad t_{ij,\alpha}^{\text{eff}} = t_{ij,\alpha} \langle b_i^\dagger b_j \rangle \quad K_{ij} = \sum_{\alpha} t_{ij,\alpha} \langle f_{i,\alpha}^\dagger f_{j,\alpha} \rangle$$

$$H_{mf} = \sum_{ij;\alpha} t_{ij,\alpha}^{\text{eff}} f_i^\dagger f_j + K_{ij} b_i^\dagger b_j + \frac{1}{2} \sum_{i,j} U_{ij} n_i n_j$$

Physics:
Phases become
U(1) gauge fields

c.f. Senthil 2008

Secondary MF

MF conditions $t_{ij,\alpha}^{\text{eff}} = t_{ij,\alpha} \langle b_i^\dagger b_j \rangle \quad K_{ij} = \sum_{\alpha} t_{ij,\alpha} \langle f_{i,\alpha}^\dagger f_{j,\alpha} \rangle$

$$H_{mf} = \sum_{ij;\alpha} t_{ij,\alpha}^{\text{eff}} f_i^\dagger f_j + K_{ij} b_i^\dagger b_j + \frac{1}{2} \sum_{i,j} U_{ij} n_i n_j$$






Boson rotor model still non-trivial: a canonical model for Bose Mott and Bose crystal transitions

For a tractable calculation we carry out a secondary MF for the boson problem, and work to quadratic order in the fluctuations around it. This is sufficient to obtain all necessary expectation values, and becomes exact in the large U limit.

Mean Field results

On-site repulsion: U nearest neighbor repulsion: $V = U/4$

next nearest neighbor repulsion: $V' = 0$




3-site and 4-site unit cell ansatz


Mean Field results

On-site repulsion: U nearest neighbor repulsion: $V = U/4$


next nearest neighbor repulsion: $V' = 1/\sqrt{3}V$




All the phases




(c) $\bar{n} = 1$




(d) $\bar{n} = 5/4$




(e) $\bar{n} = 4/3$




(f) $\bar{n} = 3/2$




(g) $\bar{n} = 5/3$




(h) $\bar{n} = 7/4$




(i) triangular metal



(j) honeycomb metal



(k) kagome metal




(l) stripe metal


?? Can we find spin liquids in any of these states?

VBS states


At mean field level, dimer (VBS) states always have the lowest energy





$$n = 1$$



$$n = 5/4$$



$$n = 4/3$$






$$= \frac{1}{\sqrt{2}} (|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle)$$

BUT...

Triangular

Dirac spin liquid states are quite energetically competitive,
and are likely to be favored after Gutzwiller projection back to physical space




Low energy theory: N=4 QED in 2+1 dim



$$\langle S^\alpha(\mathbf{x}) S^\alpha(\mathbf{y}) \rangle \sim \frac{-e^{i\mathbf{K} \cdot (\mathbf{x} - \mathbf{y})}}{|\mathbf{x} - \mathbf{y}|^{2\Delta_\phi}} + \text{h.c.} + \dots$$

$\Delta_\phi \approx 1.02$

Honeycomb



Resonance valence bond







U(1) Dirac spin liquid

Unstable to monopole proliferation

X.-Y. Song *et al.*, *Nature Commun* 2019

Kagome



U(1) Dirac spin liquid


Emergent SU(4) symmetry

Beyond MF

- Small energy differences in MF results: we need to be more precise!
- Turn to Density Matrix Renormalization Group
- Recent advances make it possible to directly study substantially large Hubbard (not Heisenberg or t-J) models, despite larger Hilbert space
- We are completely reliant upon the expertise of Hongchen Jiang (SLAC)!

Honeycomb CDW


$\bar{n} = 4/3, U = 12t, V = U/4, V' = 0, m = 25k, \text{U(1) DMRG}$



Not a spin liquid. Zig-zag spin order.

Inverse honeycomb CDW


$\bar{n} = 5/3, U = 12t, V = U/4, V' = 0, m = 25k, \text{U(1) DMRG}$



Might be a spin liquid.
No visible moments in bulk.

Weak $n=5/4$ CDW




$\bar{n} = 5/4, U = 12t, V = U/4, V' = 0, m = 25k, \text{U}(1) \text{ DMRG}$




Without NNN interaction, we see small charge order, consistent with a collinear SDW and induced secondary CDW order.


Kagome CDW

$$\bar{n} = 5/4, U = 12t, V = U/4, V' = V/\sqrt{3}, m = 8k, \text{SU}(2) \text{ DMRG}$$




Consistent with a spin liquid. Not clear what type.






Dirac QSL




- A realization of QED₃ in a spin system
- Very often found to be energetically best amongst trial wave functions.


$$\mathcal{L} = \bar{\psi}\gamma^\mu(\partial_\mu - ia_\mu)\psi + \dots$$



X.-Y. Song *et al*, 2019




Y. Iqbal *et al*, 2013



A. Wietek *et al*, 2024

YbZn₂GaO₅




Xu *et al*, 2024

Spins and QED₃

Each system has its own:

- Microscopic (exact) symmetries
- Operator dictionary
- Perturbations to CFT


X.-Y. Song *et al*, 2019




We seek some qualitative and quantitative signatures

Applied field

- Essential tuning knob available in experiment






T. Suzuki et al, 2013



M. Jaime et al, 2006

- Field couples via Zeeman term $-HS^z$
- Non-zero dM/dH usually indicates symmetry breaking

Proof of principle




R. Coldea et al, 2010

Applied field

$$H = H_{\text{QED3}} - h\psi^\dagger \sigma^z \psi$$

- Field couples to conserved SU(2) charge $M = S^z$
- SU(2)_{spin} is a subgroup of SU(N) symmetry of QED3
- Conformal QED3 also has emergent U(1)_{flux} conservation symmetry



States labeled by M and Φ

$$E = E_0(M, \Phi) - hM$$

Applied field

$$E = E_0(M, \Phi) - hM$$

- Key question: what value of Φ minimizes E ?


[Y. Ran *et al*, 2009]: $\Phi = \pm 2\phi_0$

n.b. sign
breaks TRS




LL state

- Ran *et al*: LL state is a magnetically ordered state!
 - It breaks TRS
 - It has a fermion gap, so monopole-monopole correlations are long-ranged
 - Response to probe gauge field A_{spin}^{μ} shows that $U(1)_{\text{spin}}$ symmetry is spontaneously broken



LL state

- Ran *et al*: LL state is a magnetically ordered state!
 - It breaks TRS
 - It has a fermion gap, so monopole-monopole correlations are long-ranged
 - Response to probe gauge field A_{spin}^{μ} shows that $U(1)_{\text{spin}}$ symmetry is spontaneously broken




Can we still see signs
of QED3 in the
ordered phase?

Spectroscopy

- Ran *et al* analysis of ordered phase is entirely based on energies $\ll \hbar\omega_c$
- QED3 theory should be valid for any small energy, including multiples of $\hbar\omega_c$
- Project: seek signs of QED3 structure on these scales in observables.

Dynamical spin correlations



LL transitions:
"magneto-excitons"

$$\chi_{\pm}(k, \omega) = \text{bubble diagram} + \text{bubble with wavy line} + \text{bubble with wavy line} + \text{bubble with wavy line} + \dots$$

The equation shows a series of diagrams for the dynamical spin correlation function $\chi_{\pm}(k, \omega)$. Each diagram consists of a bubble (two vertices connected by two lines) with an upward arrow $\uparrow$ above and a downward arrow $\downarrow$ below. The first diagram is a simple bubble. The subsequent three diagrams show a wavy line (representing a magneto-exciton) inserted into the bubble, with the wavy line connecting the two vertices. The series continues with an ellipsis $\dots$.

"Excitonic" corrections will give
dispersion, broadening



A fully worked example


A. Keselman *et al* (2020)

$$H = \sum_i \left[J_1 \vec{S}_i \cdot \vec{S}_{i+1} + J_2 \vec{S}_i \cdot \vec{S}_{i+2} - B S_i^z \right]$$

- Bosonization and reffermionization:

$$H_0 = v \int dx \left(\psi_R^\dagger (-i\partial_x) \psi_R + \psi_L^\dagger (i\partial_x) \psi_L \right)$$

$$V = -g \int dx \vec{J}_R \cdot \vec{J}_L = -g \int dx \left[J_R^z J_L^z + \frac{1}{2} J_R^+ J_L^- + \frac{1}{2} J_R^- J_L^+ \right]$$



$$\vec{S}_i \sim \vec{J}_R(x_i) + \vec{J}_L(x_i) + (-1)^i \vec{N}(x_i)$$

$$\vec{J}_{R/L} = \frac{1}{2} \psi_{R/L}^\dagger \vec{\sigma} \psi_{R/L}$$

Small k spin correlations ~ fermion bilinear




A fully worked example

MPS methods: DMRG+TEBD

Near QCP


$$g \approx 0$$



well-approximated by free fermion spin correlations (at small k)

NN chain

$$B/J_1 = 1$$



Excitonic corrections evident due to interaction g (at small k)




A fully worked example

MPS methods: DMRG+TEBD

Near QCP


$$g \approx 0$$





well-approximated by free
fermion spin correlations (at
small k)

NN chain

$$B/J_1 = 1$$



Excitonic corrections evident
due to interaction g (at small k)




The QED3 case

$$\chi_{\pm}(k, \omega) = \text{[Feynman diagram series]}$$

The diagram series shows a bubble diagram with external arrows, followed by a sum of diagrams with internal wavy lines and dots, representing a perturbative expansion.

$$\begin{aligned} \text{Im}[\Pi^{+-}(\nu + i\eta, \mathbf{k})] &\sim \pi \times \frac{e^{-|\mathbf{k}|^2 \ell^2/2}}{2\pi \ell^2} \left[\delta_{\eta}(\nu - 2h) + \sum_{n=1}^{\infty} \frac{1}{n!} \left(\frac{\ell^2 |\mathbf{k}|^2}{2} \right)^n \delta_{\eta}(\nu - 2h - E_n) \right. \\ &+ \frac{1}{4} \sum_{n=1, m=1}^{\infty} (-1)^{n+m} [\delta_{\eta}(\nu - (E_m + E_n + 2h)) - \delta_{\eta}(\nu - (-E_m - E_n + 2h))] \times \{ \mathcal{L}_{0,0}^{n,m}(\ell^2 |\mathbf{k}|^2/2) + \mathcal{L}_{1,1}^{n,m}(\ell^2 |\mathbf{k}|^2/2) \\ &\quad \left. - 2 \frac{E_n}{E_m} \mathcal{L}_{0,1}^{n,m}(\ell^2 \mathbf{k}^2/2) \} \right] \end{aligned}$$

ω/h



Will be interesting to see how this structure is affected by excitonic corrections

Issues

- Full treatment of gauge fluctuations: lattice gauge simulation? (No sign problem!)
- Some spin correlations are represented by monopole operators: how to calculate these contributions?
- Role of perturbations to the CFT Hamiltonian

Thank you

