




Quantum kinetics, non-linear response, and band geometry

Collaborators



Lucile Savary
FACTS



Takamori Park
UCSB -> Princeton



Xiaoyang Huang
CU Boulder -> PI



Wen Wang
UCSB



Jia-Xin Zhang
FACTS




Dongning (Donna) Sheng
CSUN

Outline

- Bands and quantum geometry
- Quantum geometric nesting in flat bands
- Doing semiclassics correctly in dispersive bands with quantum geometry
- A quick advertisement for recent work on topological superconductivity

Bands



$|\psi_{nk}\rangle$

Geometry


$$\epsilon_n(k) \rightarrow \mathbf{v}_n(k)$$

Energetics

$\Omega_{n,\alpha\beta}(k)$ Berry curvature

$g_{n,\alpha\beta}(k)$ Quantum metric

Quantum geometry




Phase of Bloch state varies with k

Berry phase around a loop is given by the **Berry curvature**.

$$\Omega_{n,\alpha\beta}(k)$$

Quantum geometry




Wavefunction of Bloch state varies with k

How fast they vary is described by **quantum metric**

$$|\langle \psi_{nk} | \psi_{nk+q} \rangle|^2 \sim 1 - g_{\mu\nu}(k) q_\mu q_\nu$$

Quantum geometry




Wavefunction of Bloch state varies with k

How fast they vary is described by **quantum metric**

$$|\langle \psi_{nk} | \psi_{nk+q} \rangle|^2 \sim 1 - g_{\mu\nu}(k) q_\mu q_\nu$$

Quantum geometry



Example: Haldane model

Wavefunction of Bloch state varies with k

How fast they vary is described by **quantum metric**


$$|\langle \psi_{nk} | \psi_{nk+q} \rangle|^2 \sim 1 - g_{\mu\nu}(k) q_{\mu} q_{\nu}$$

Ellipses show constant “distance”

Quantum geometric nesting




Wen Wang
UCSB



Jia-Xin Zhang
FACTS

Bands



$|\psi_{nk}\rangle$

Geometry


$$\epsilon_n(k) \rightarrow \mathbf{v}_n(k)$$

Energetics

$\Omega_{n,\alpha\beta}(k)$ Berry curvature

$g_{n,\alpha\beta}(k)$ Quantum metric

Flat bands



$|\psi_{nk}\rangle$


Geometry

In the flat band limit, there is *only* geometry

$\Omega_{n,\alpha\beta}(k)$ Berry curvature

$g_{n,\alpha\beta}(k)$ Quantum metric

Flat bands



In the flat band limit, there is *only* geometry

Interaction $U \ll \Delta$

How does geometry determine the instabilities?

$\Omega_{n,\alpha\beta}(k)$ Berry curvature



$g_{n,\alpha\beta}(k)$ Quantum metric

Flat bands

Generally a result of interference in electron motion, which can be induced by frustration or by magnetic flux.



Landau levels

Frustrated hopping




Flat bands

moiré



Bistritzer+Macdonald, 2011


Graphene




F. Wu *et al*, 2019

TMDs

Correlation physics




FQHE



Cao *et al*, 2018

Superconductivity and correlated insulators in TBG



H. Park *et al*, 2023


FQAHE in twisted MoTe2

So many different orders

- Ferromagnetism
- Antiferromagnetism
- Intervalley coherence
- Fractional quantum Hall effect
- Stripes
- Charge density wave
- Loop currents
- Superconductivity (s-wave, p-wave, d-wave, FFLO...)


Nesting

Within a *dispersive* band, Fermi surface structure controls instabilities



Quasi-1d


" $2k_F$ " CDW/SDW



Square lattice Hubbard

(π, π) antiferromagnet

Susceptibility



Restricts to near Fermi surface

$$\chi(Q, \omega) \sim \sum_k \frac{n_F(\epsilon_{k+Q}) - n_F(\epsilon_k)}{\omega - \epsilon_k + \epsilon_{k+Q}}$$

Enhanced susceptibility
when this vanishes for
extended range of k

What happens when the band is flat?? No Fermi surface!

Quantum geometric nesting

Z. Han *et al*, 2024


For flat band (or bands) at the Fermi energy, susceptibility is rendered finite only by thermal fluctuations. But there is still structure.

$$\chi_{\mathbf{Q}} = \frac{1}{T} \sum_{\mathbf{k}} \text{Tr} [\mathcal{O}^\dagger(\mathbf{k}) P(\mathbf{k} + \mathbf{Q}) \mathcal{O}(\mathbf{k}) P(\mathbf{k})] .$$

Order parameter

Projector

$$P(\mathbf{k}) = \sum_{n \in \text{FB}} |\psi_{n\mathbf{k}}\rangle \langle \psi_{n\mathbf{k}}|$$




Quantum geometric nesting


Bloch vector representation

e.g. 2-band

$$P(\mathbf{k}) = \frac{1}{2} + \frac{1}{2} \mathbf{b}(\mathbf{k}) \cdot \boldsymbol{\sigma}$$



$$\mathcal{O}(\mathbf{k}) = o(\mathbf{k}) + \mathbf{o}(\mathbf{k}) \cdot \boldsymbol{\sigma}$$




Quantum geometric nesting

Bloch vector representation


N bands, N_L flat

$$P(\mathbf{k}) = \frac{N_L}{N} + \frac{1}{2} \mathbf{b}(\mathbf{k}) \cdot \boldsymbol{\lambda}$$

$$\boldsymbol{\lambda} \in \text{SU}(N)$$



$$\mathcal{O}(\mathbf{k}) = o(\mathbf{k}) + \mathbf{o}(\mathbf{k}) \cdot \boldsymbol{\lambda}$$




Quantum geometric nesting

$$\chi(\mathbf{Q}) = \sum_{\mathbf{k}} \frac{\text{Tr}[\mathcal{O}^\dagger(\mathbf{k})\mathcal{O}(\mathbf{k})]}{T} \left(\frac{N_L^2}{N} + \frac{1}{2} \tilde{\mathbf{b}}_o(\mathbf{k} + \mathbf{Q}) \cdot \mathbf{b}(\mathbf{k}) \right),$$

Relates quantum
geometry at $\mathbf{k}$ and $\mathbf{k}+\mathbf{Q}$

$$\tilde{\mathbf{b}}_o(\mathbf{k} + \mathbf{Q}) \equiv \mathbf{b}(\mathbf{k} + \mathbf{Q}) - N [\hat{\mathbf{o}}(\mathbf{k}) \times \mathbf{b}(\mathbf{k} + \mathbf{Q}) \times \hat{\mathbf{o}}(\mathbf{k})]$$



QGN condition:



$$\tilde{\mathbf{b}}_o(\mathbf{k} + \mathbf{Q}) \parallel \mathbf{b}(\mathbf{k}), \quad \forall \mathbf{k} \in \text{BZ}$$

Example



JS Hofmann et al, 2022,2023

$$H_0 = t \sum_{\mathbf{k}} c_{\mathbf{k}}^{\dagger} \begin{pmatrix} -\mu & -ie^{i\alpha_{\mathbf{k}}^{\uparrow}} & 0 & 0 \\ ie^{-i\alpha_{\mathbf{k}}^{\uparrow}} & -\mu & 0 & 0 \\ 0 & 0 & -\mu & ie^{-i\alpha_{\mathbf{k}}^{\downarrow}} \\ 0 & 0 & -ie^{i\alpha_{\mathbf{k}}^{\downarrow}} & -\mu \end{pmatrix} c_{\mathbf{k}}$$

$$\alpha_{\mathbf{k}}^{\sigma} = \eta_{\sigma} (\cos k_x + \cos k_y)$$



$b(\mathbf{k})$






$\tilde{b}_N(\mathbf{k})$

Example

JS Hofmann et al, 2022,2023

$$H_0 = t \sum_{\mathbf{k}} c_{\mathbf{k}}^{\dagger} \begin{pmatrix} -\mu & -ie^{i\alpha_{\mathbf{k}}^{\uparrow}} & 0 & 0 \\ ie^{-i\alpha_{\mathbf{k}}^{\uparrow}} & -\mu & 0 & 0 \\ 0 & 0 & -\mu & ie^{-i\alpha_{\mathbf{k}}^{\downarrow}} \\ 0 & 0 & -ie^{i\alpha_{\mathbf{k}}^{\downarrow}} & -\mu \end{pmatrix} c_{\mathbf{k}}$$

$$\alpha_{\mathbf{k}}^{\sigma} = \eta_{\sigma} (\cos k_x + \cos k_y)$$





Example

JS Hofmann et al, 2022,2023

$$H_0 = t \sum_{\mathbf{k}} c_{\mathbf{k}}^\dagger \begin{pmatrix} -\mu & -ie^{i\alpha_k^\uparrow} & 0 & 0 \\ ie^{-i\alpha_k^\uparrow} & -\mu & 0 & 0 \\ 0 & 0 & -\mu & ie^{-i\alpha_k^\downarrow} \\ 0 & 0 & -ie^{i\alpha_k^\downarrow} & -\mu \end{pmatrix} c_{\mathbf{k}}$$

$$\alpha_k^\sigma = \eta_\sigma (\cos k_x + \cos k_y)$$



$b(\mathbf{k})$


QGN!

QMC Simulation

Include Hubbard interaction $H = H_0 + U \sum_i \left(n_{i,A}^2 + n_{i,B}^2 \right)$

Use dQMC for 8x8 lattice



$$\mathbf{Q} = (\pi, \pi)$$



QMC Simulation

Include Hubbard interaction $H = H_0 + U \sum_i \left(n_{i,A}^2 + n_{i,B}^2 \right)$

Use dQMC for 8x8 lattice



Superconductivity


With attractive interactions in the flat band, can get exotic superconductivity

$$\chi_Q^{\text{pp}} = \frac{1}{4T} \frac{|\nu|}{\text{arctanh}|\nu|} \sum_{\mathbf{k}} o_0^2(\mathbf{k}) \left(Nn^2 + \frac{1}{2} \mathbf{b}^R(\mathbf{k} + \mathbf{Q}) \cdot \mathbf{b}(-\mathbf{k}) \right)$$


Similar nesting conditions

$$\mathbf{b}^R(\mathbf{k} + \mathbf{Q}) \parallel \mathbf{b}(-\mathbf{k})$$


$$\mathbf{b}^R\left(\frac{\mathbf{Q}}{2} + \frac{\mathbf{k}}{2}\right) \parallel \mathbf{b}\left(\frac{\mathbf{Q}}{2} - \frac{\mathbf{k}}{2}\right)$$




Superconducting QGN



QMC




$$\eta_{\sigma} = 0.75\sigma$$



Summary

- We developed an algebraic approach to quantum geometric nesting
- It allows intuitive study of ordering instabilities of flat bands
- QMC validates the approach for a simple model system
- We also obtain relations between a “high temperature stiffness” and a generalized quantum metric (for both particle/hole and superconducting orders at any $\mathbf{Q}$)
- ?? Is there a way to include *non-symmetry breaking “orders”* like FQAHE?

Kinetic theory




Takamori Park
UCSB -> Princeton



Xiaoyang Huang
CU Boulder -> PI


Bands



$|\psi_{nk}\rangle$

Geometry

$\epsilon_n(k)$



$\mathbf{v}_n(k)$

Energetics

$$\sigma_{xx} \sim \langle v_n^x v_n^x \rangle$$

$\Omega_{n,\alpha\beta}(k)$

Berry curvature

$g_{n,\alpha\beta}(k)$

Quantum metric

$$\sigma_{xy} \sim \langle \Omega \rangle$$

Souza, Wilkens, Martin 2000
Various suggestions

Role in interacting systems?

Linear response

A host of “conventional” (but still complex) transport coefficients

$$\begin{aligned}\mathbf{j} &= \mathbf{L}^{11}\boldsymbol{\varepsilon} + \mathbf{L}^{12}(-\nabla T), \\ \mathbf{j}^q &= \mathbf{L}^{21}\boldsymbol{\varepsilon} + \mathbf{L}^{22}(-\nabla T),\end{aligned}$$

Ashcroft+Mermin

And some originating from Berry curvature

Anomalous Hall effect

$$\mathbf{j}_e^{ah} = e^2 \sum_n \int \frac{d^d \mathbf{k}}{(2\pi)^d} n_F(\omega_n(\mathbf{k})) \boldsymbol{\Omega}_n(\mathbf{k}) \times \mathbf{E}$$

Karplus+Luttinger, 1954!

Gyromagnetic effect

$$\alpha_{ij}^{\text{GME}} = \frac{i\omega\tau e}{1 - i\omega\tau} \sum_n \int [d\mathbf{k}] (\partial f / \partial \varepsilon_{\mathbf{k}n}) v_{\mathbf{k}n,i} m_{\mathbf{k}n,j}$$

Ma+Pesin 2015; Song et al 2016

Chiral anomaly...


Non-linear response

Non-linear Hall effect

$$j_a^0 = \chi_{abc} \mathcal{E}_b \mathcal{E}_c^*, \quad j_a^{2\omega} = \chi_{abc} \mathcal{E}_b \mathcal{E}_c,$$

$$\chi_{abc} = -\varepsilon_{adc} \frac{e^3 \tau}{2(1 + i\omega\tau)} \int_k f_0(\partial_b \Omega_d)$$

Sodemann+Fu, 2015



Q. Ma *et al*, 2019 (WTe2)

Non-linear response

Quantum metric effects


$$\sigma_{a,bc}^{\text{BCPD}} = \frac{e^3}{\hbar} \sum_{m,p,k} \int [dk] f_m [\partial_a \tilde{\mathcal{G}}_{mp}^{bc} + \partial_b \tilde{\mathcal{G}}_{mp}^{ac} + \partial_c \tilde{\mathcal{G}}_{mp}^{ab}].$$

“Band-resolved quantum metric dipole”


K. Das et al, 2023

D. Kaplan et al, 2024

MnBi₂Te₄



N. Wang et al, 2023



A. Gao et al, 2023

Quadrupole response

$$j_{\text{geom.}}^{\mu}(\mathbf{r}) = -\frac{Q^2}{2\hbar} \int \frac{d^D \mathbf{K}}{(2\pi)^D} f_0(\mathbf{K}) E_{\nu\lambda}^{(0)} \frac{\partial g^{\nu\lambda}(\mathbf{K})}{\partial K_{\mu}},$$

Quantum metric dipole

M. Lapa + T. Hughes, 2019


$$E_{\mu\nu} = \partial E_{\mu} / \partial x_{\nu}$$

Semiclassical approach

Two approximations, done separately

c.f. Great talk from
Jenny Coulter

Band structure




$$|\psi_{nk}\rangle$$

Single particle
equations of motion

$$\begin{aligned}\frac{d\mathbf{x}}{dt} &= \mathbf{v}_n = \nabla_{\mathbf{k}} \tilde{\epsilon}_{n\mathbf{k}} - \frac{d\mathbf{k}}{dt} \times \boldsymbol{\Omega}_{n\mathbf{k}}, \\ \frac{d\mathbf{k}}{dt} &= \mathbf{F}_n = -e\mathbf{E} - e\frac{d\mathbf{x}}{dt} \times \mathbf{B}.\end{aligned}$$

Boltzmann equation



$$\partial_t f_n + \frac{dx_\mu}{dt} \frac{\partial f_n}{\partial x_\mu} + \frac{dk_\mu}{dt} \frac{\partial f_n}{\partial k_\mu} = \mathcal{C}[f]$$

$$[x, k] = i\hbar$$

$$\begin{pmatrix} \rho_1 & \rho_{12} \\ \rho_{21} & \rho_2 \end{pmatrix} \rightarrow \begin{pmatrix} f_1 & 0 \\ 0 & f_2 \end{pmatrix}$$

What we did


Systematic exact (order by order in spatial gradients) derivation of kinetics from the density matrix alone.

Just a hint of the (rather technical) method:

$$i\hbar\partial_t\rho = [H, \rho]$$

Matrix in *both* band and position/momentum space


$$\rho_{n,k;n',k'}$$



Wigner transform

$$i\hbar\partial_t\mathbf{F} = [\mathbf{H}, \mathbf{F}]_\star$$

$$F_{n,n'}(k, x)$$



Moyal diagonalization

QKE

$$f_n(k, x)$$


Kinetic equation

$$\partial_t f_n + \partial_\alpha \mathcal{J}_\alpha = 0$$

Just continuity equation for density in a band

c.f. simple convection

$$\mathcal{J}_\alpha = f_n v_\alpha$$



Incompressible flow

Boltzmann equation

Here, to second order:

$$\begin{aligned} \mathcal{J}_\alpha = & \varepsilon_{\alpha\beta} \operatorname{tr} \left[f \left(\partial_\beta h + \hbar (\varepsilon_{\sigma\lambda} \partial_\sigma h \Omega_{\lambda\beta} - \Omega_{t\beta}) \left(1 - \frac{\hbar}{2} \Omega_{\mu\nu} \right) + \frac{\hbar^2}{2} \varepsilon_{\sigma\lambda} \varepsilon_{\mu\nu} \partial_{\sigma\mu}^2 h \partial_\beta g_{\nu\lambda} \right) \right] \\ & + \hbar^2 \varepsilon_{\alpha\beta} \varepsilon_{\sigma\lambda} \partial_\sigma \operatorname{tr} \left[f \left(\frac{1}{2} \varepsilon_{\mu\nu} \partial_\mu h \partial_\nu g_{\beta\lambda} - \frac{1}{2} \partial_t g_{\beta\lambda} + \varepsilon_{\mu\nu} \partial_{\mu\beta}^2 h g_{\nu\lambda} \right) \right] - \frac{\hbar^2}{24} \varepsilon_{\alpha\beta} \varepsilon_{\sigma\lambda} \varepsilon_{\mu\nu} \partial_{\sigma\mu}^2 \operatorname{tr} [f \partial_{\beta\lambda\nu}^3 h] \end{aligned}$$


Kinetic equation

$$\partial_t f_n + \partial_\alpha \mathcal{J}_\alpha = 0$$

Just continuity equation for density in a band

c.f. simple convection

$$\mathcal{J}_\alpha = f_n v_\alpha$$



Incompressible flow

Boltzmann equation

Here, to second order:

$$\begin{aligned} \mathcal{J}_\alpha = & \varepsilon_{\alpha\beta} \operatorname{tr} \left[f \left(\partial_\beta h + \hbar (\varepsilon_{\sigma\lambda} \partial_\sigma h \Omega_{\lambda\beta} - \Omega_{t\beta}) \left(1 - \frac{\hbar}{2} \Omega_{\mu\nu} \right) + \frac{\hbar^2}{2} \varepsilon_{\sigma\lambda} \varepsilon_{\mu\nu} \partial_{\sigma\mu}^2 h \partial_\beta g_{\nu\lambda} \right) \right] \\ & + \hbar^2 \varepsilon_{\alpha\beta} \varepsilon_{\sigma\lambda} \partial_\sigma \operatorname{tr} \left[f \left(\frac{1}{2} \varepsilon_{\mu\nu} \partial_\mu h \partial_\nu g_{\beta\lambda} - \frac{1}{2} \partial_t g_{\beta\lambda} + \varepsilon_{\mu\nu} \partial_{\mu\beta}^2 h g_{\nu\lambda} \right) \right] - \frac{\hbar^2}{24} \varepsilon_{\alpha\beta} \varepsilon_{\sigma\lambda} \varepsilon_{\mu\nu} \partial_{\sigma\mu}^2 \operatorname{tr} [f \partial_{\beta\lambda\nu}^3 h] \end{aligned}$$

quantum metric $g_{\alpha\beta} = \frac{1}{2} \operatorname{diag} (\Lambda_\alpha \Lambda_\beta + \Lambda_\beta \Lambda_\alpha) - A_\alpha A_\beta = \frac{1}{2} \operatorname{tr} (\partial_\alpha P_n \partial_\beta P_n)$

Kinetic equation

$$\begin{aligned} \mathcal{J}_\alpha = & \varepsilon_{\alpha\beta} \operatorname{tr} \left[f \left(\partial_\beta h + \hbar (\varepsilon_{\sigma\lambda} \partial_\sigma h \Omega_{\lambda\beta} - \Omega_{t\beta}) \left(1 - \frac{\hbar}{2} \Omega_{\mu\nu} \right) + \frac{\hbar^2}{2} \varepsilon_{\sigma\lambda} \varepsilon_{\mu\nu} \partial_{\sigma\mu}^2 h \partial_\beta g_{\nu\lambda} \right) \right] \\ & + \hbar^2 \varepsilon_{\alpha\beta} \varepsilon_{\sigma\lambda} \partial_\sigma \operatorname{tr} \left[f \left(\frac{1}{2} \varepsilon_{\mu\nu} \partial_\mu h \partial_\nu g_{\beta\lambda} - \frac{1}{2} \partial_t g_{\beta\lambda} + \varepsilon_{\mu\nu} \partial_{\mu\beta}^2 h g_{\nu\lambda} \right) \right] - \frac{\hbar^2}{24} \varepsilon_{\alpha\beta} \varepsilon_{\sigma\lambda} \varepsilon_{\mu\nu} \partial_{\sigma\mu}^2 \operatorname{tr} [f \partial_{\beta\lambda\nu}^3 h] \end{aligned}$$

- General and exact up to second order in semiclassical expansion
- Explicitly contains real space and momentum space, and capable of describing inhomogeneous systems. Magnetic field can be included by modifying $\epsilon_{\alpha\beta}$
- Quantum geometry of bands appears explicitly, takes local band Hamiltonian as input
- **Only band-intrinsic quantities appear**: but these are bands renormalized by quantum corrections.
- Ready to attack all sorts of problems!

Example: separable problem

$$H(x, p) = H_0(p) - eV(x)I_N$$

Band form Diagonal potential

Kinetic equation (relaxation time approx)

$$\begin{aligned} \partial_t f = & -\partial_{x_i} \left[f \left(v_i + \frac{e}{\hbar} \Omega_{ij} \partial_{x_j} V + e \mathsf{T}_{ij} \partial_{tx_j}^2 V + \frac{e^2}{\hbar} \left(\partial_{k_j} \mathsf{T}_{il} - \frac{1}{2} \partial_{k_i} \mathsf{T}_{jl} \right) \partial_{x_j} V \partial_{x_l} V + \frac{e}{\hbar} \left(\hbar v_j \mathsf{T}_{il} - \frac{1}{2} \partial_{k_i} \mathsf{g}_{jl} \right) \partial_{x_j x_l}^2 V \right) \right] \\ & - \frac{e}{\hbar} \partial_{k_i} f \partial_{x_i} V + \frac{e}{2\hbar} \partial_{x_i x_l}^2 (f \partial_{k_j} \mathsf{g}_{il} \partial_{x_j} V) - \frac{e}{\hbar} \partial_{k_i x_j}^2 (f \mathsf{g}_{jl} \partial_{x_i x_l}^2 V) + \frac{1}{24\hbar} \partial_{x_i x_j x_l}^3 f \partial_{k_i k_j k_l}^3 \varepsilon \\ & + \frac{e}{24\hbar} \partial_{k_i k_j k_l}^3 f \partial_{x_i x_j x_l}^3 V + \mathcal{I}[f] + \mathcal{O}(\partial_x^4). \end{aligned} \quad (3.29)$$

$$T_{n;\mu\nu} = - \sum_{m \neq n} \frac{\langle \partial_{p_\mu} u_n | u_m \rangle \langle u_m | \partial_{p_\nu} u_n \rangle + (\mu \leftrightarrow \nu)}{\epsilon_n - \epsilon_m} \quad \begin{array}{l} \text{"band-normalized quantum metric"} \\ \text{(not a purely geometric quantity)} \end{array}$$

Results

- Reproduced
 - Non-linear Hall effect
 - Non-linear band-resolved quantum metric dipole contributions
 - Quadrupole response
 - New results in collision less regime
 - e.g. polarization function
- } General form agrees with literature but we found errors

$$\hat{C}^{(2)}(\omega, q) = -ie^{-1}\hbar \int_k \frac{\delta \hat{f}_{e=1}(q, k, \omega)}{\delta \hat{V}(q, \omega)} \approx -i \int_k \left[\frac{\hbar^2 n'_F \vec{v} \cdot \vec{q} (1 - \mathbf{g}_{ij}(k) q_i q_j)}{\hbar \omega - \varepsilon(k + q/2) + \varepsilon(k - q/2)} + \hbar n_F \mathbf{t}_{ij}(k) q_i q_j \right]$$

Summary

$$\begin{aligned} \mathcal{J}_\alpha = & \varepsilon_{\alpha\beta} \operatorname{tr} \left[f \left(\partial_\beta h + \hbar (\varepsilon_{\sigma\lambda} \partial_\sigma h \Omega_{\lambda\beta} - \Omega_{t\beta}) \left(1 - \frac{\hbar}{2} \Omega_{\mu\nu} \right) + \frac{\hbar^2}{2} \varepsilon_{\sigma\lambda} \varepsilon_{\mu\nu} \partial_{\sigma\mu}^2 h \partial_\beta g_{\nu\lambda} \right) \right] \\ & + \hbar^2 \varepsilon_{\alpha\beta} \varepsilon_{\sigma\lambda} \partial_\sigma \operatorname{tr} \left[f \left(\frac{1}{2} \varepsilon_{\mu\nu} \partial_\mu h \partial_\nu g_{\beta\lambda} - \frac{1}{2} \partial_t g_{\beta\lambda} + \varepsilon_{\mu\nu} \partial_{\mu\beta}^2 h g_{\nu\lambda} \right) \right] - \frac{\hbar^2}{24} \varepsilon_{\alpha\beta} \varepsilon_{\sigma\lambda} \varepsilon_{\mu\nu} \partial_{\sigma\mu}^2 \operatorname{tr} [f \partial_{\beta\lambda\nu}^3 h] \end{aligned}$$

- We found a procedure to obtain semiclassical electron dynamics organized by spatial gradients
- We found a general result for the phase space current to *second* order in gradients, **capable of handling inhomogeneous systems and magnetic fields**
- Prior results for linear and non-linear response are recovered, and in some cases corrected
- New results for collision less regime
- Still needed: a scattering/interactions theory with the same validity


Brief update: anyon superconductivity?

Chiral Spin Liquid and Quantum Phase Transition in the Triangular Lattice Hofstadter-Hubbard Model

Stefan Divic,^{1,2} Tomohiro Soejima,³ Valentin Crépel,² Michael P. Zaletel,^{1,4} and Andrew Millis^{2,5}

Anyon Superconductivity from Topological Criticality in a Hofstadter-Hubbard Model



Stefan Divic,¹ Valentin Crépel,² Tomohiro Soejima (副島智大),³ Xue-Yang Song,⁴ Andrew J. Millis,^{2,5} Michael P. Zaletel,^{1,6} and Ashvin Vishwanath³



Charge gap must vanish at this QCP, but 1-e gap does not.

Strong indication that paired state arises on doping.

Our study




Strong pairing correlations

Robust superconductivity upon doping chiral spin liquid and Chern insulators in a Hubbard-Hofstadter model

Clemens Kuhlenkamp,^{1,*} Stefan Divic,^{2,3,*} Michael P. Zaletel,^{3,4}
Tomohiro Soejima (副島智大),¹ and Ashvin Vishwanath¹


c.f.

Topology: spin pumping



DQMC


Sign problem, but manageable



BCS or not?




All in the superconducting phase, but we can see the evolution

Spin $S(\mathbf{k})$



Occupation number $n(\mathbf{k})$

Thanks



arXiv: 2509.02757

arXiv: 2504.03882

$$\mathcal{J}_\alpha = \varepsilon_{\alpha\beta} \text{tr} \left[f \left(\partial_\beta h + \hbar (\varepsilon_{\sigma\lambda} \partial_\sigma h \Omega_{\lambda\beta} - \Omega_{t\beta}) \left(1 - \frac{\hbar}{2} \Omega_{\mu\nu} \right) + \frac{\hbar^2}{2} \varepsilon_{\sigma\lambda} \varepsilon_{\mu\nu} \partial_\sigma^2 h \partial_\beta g_{\nu\lambda} \right) \right] \\ + \hbar^2 \varepsilon_{\alpha\beta} \varepsilon_{\sigma\lambda} \partial_\sigma \text{tr} \left[f \left(\frac{1}{2} \varepsilon_{\mu\nu} \partial_\mu h \partial_\nu g_{\beta\lambda} - \frac{1}{2} \partial_t g_{\beta\lambda} + \varepsilon_{\mu\nu} \partial_\mu^2 h g_{\nu\lambda} \right) \right] - \frac{\hbar^2}{24} \varepsilon_{\alpha\beta} \varepsilon_{\sigma\lambda} \varepsilon_{\mu\nu} \partial_\sigma^2 \text{tr} [f \partial_{\beta\lambda\nu}^3 h]$$

arXiv: 2504.10447

